



Movement Disorder Agents

WA.PHAR.51 Movement Disorder Agents

Background:

Huntington’s disease is a genetic, progressive, neurodegenerative disorder clinically characterized by a triad of motor, cognitive and psychiatric symptoms. Motor features include involuntary jerking or writhing movements (chorea) and voluntary movements; reduced manual dexterity, slurred speech, swallowing difficulties, balance problems and falls. Signs and symptoms develop in their 30s or 40s for most people but the disease may emerge earlier or later in life.

Tardive dyskinesia is a neurological disorder caused by the long-term use of neuroleptic drugs, or anti-psychotic medications that result in involuntary, repetitive body movements. The symptoms may include grimacing, sticking out the tongue, smacking of the lips, rapid jerking movements or slow writhing movements.

Medical necessity

Drug	Medical Necessity
deutetrabenzine (AUSTEDO™) tetrabenzazine (XENAZINE®)	deutetrabenzine and tetrabenzazine may be considered medically necessary for the diagnosis of chorea associated with Huntington’s disease or tardive dyskinesia.
valbenazine (INGREZZA®)	Valbenazine may be considered medically necessary for the diagnosis of tardive dyskinesia

Clinical policy:

Drug	Clinical Criteria (Initial Approval)
deutetrabenzine (AUSTEDO™) tetrabenzazine (XENAZINE®)	<ol style="list-style-type: none"> 1. Diagnosis of ONE of the following: <ol style="list-style-type: none"> a. Chorea associated with Huntington’s disease b. Tardive dyskinesia 2. Greater than or equal to (\geq) 18 years of age 3. Not used in combination with another vesicular monoamine transporter 2 (VMAT2) -inhibitor (e.g. tetrabenzazine, deutetrabenzine, valbenazine) 4. NONE of the following: <ol style="list-style-type: none"> a. Hepatic impairment b. Concurrent use or recent discontinuation of MAOIs or reserpine 5. Prescribed by or in consultation with a psychiatrist or neurologist 6. For deutetrabenzine only: <ol style="list-style-type: none"> a. Less than or equal to (\leq) 48mg per day 7. For tetrabenzazine only <ol style="list-style-type: none"> a. ONE of the following dose limits:

	<ol style="list-style-type: none"> i. Diagnosis of Chorea associated with Huntington’s disease less than or equal to (\leq) 50mg per day <ol style="list-style-type: none"> 1) For doses, greater than 50mg per day genotyping for CYP2D6 is required to determine if client is an intermediate or extensive metabolizer. ii. Diagnosis of tardive dyskinesia less than or equal to (\leq) 200mg per day <p>Approve for 12 months</p>
valbenazine (INGREZZA [®])	<ol style="list-style-type: none"> 1. Diagnosis of tardive dyskinesia 2. Greater than or equal to (\geq) 18 years of age 3. Not used in combination with another vesicular monoamine transporter 2 (VMAT2) -inhibitor (e.g. tetrabenazine, deutetrabenazine, valbenazine) 4. Less than or equal to (\leq) 80mg per day 5. NONE of the following: <ol style="list-style-type: none"> a. History of congenital long QT syndrome or with arrhythmias associated with a prolonged QT interval b. History of severe renal impairment c. Concomitant use with MAOIs. 6. Prescribed by or in consultation with a psychiatrist or neurologist <p>Approve for 12 months</p> <p>Criteria (Reauthorization)</p> <p>Documentation of positive clinical response</p> <p>Approve for 12 months</p>

Dosage and quantity limits

Drug Name	Dose and Quantity Limits
deutetrabenzine (AUSTEDO [™])	48mg per day;
tetrabenazine (XENAZINE [®])	HD Chorea: <ul style="list-style-type: none"> • Extensive/intermediate metabolizers: 100 mg/day • Poor metabolizers: 50 mg/day TD: 200mg per day
valbenazine (INGREZZA [®])	80mg per day;

References

1. Austedo Prescribing Information. Teva Pharmaceuticals Inc. August 2017.
2. Xenazine [Prescribing Information]. Lundbeck Inc.: Deerfield, IL. June 2015.
3. Ingrezza [Prescribing Information]. Neurocrine Pharmaceuticals; San Diego, CA. April 2017.
4. Armstrong MJ, Miyasaki JM. Evidence-based guideline: Pharmacologic treatment of chorea in Huntington disease: Report of the Guideline Development Subcommittee of the American Academy of Neurology. Neurology. 2012 August.

5. Claassen DO, Carroll B, De Boer LM, et al. Indirect tolerability comparison of deutetrabenazine and tetrabenazine for Huntington disease. *J Clin Mov Disord*. 2017. 4:3.
6. Geschwind MD, Paras N. Deutetrabenazine for treatment of chorea in Huntington disease. *JAMA*. 2016; 316(1):33-34.
7. Huntington Study Group. Effect of deutetrabenazine on chorea among patients with Huntington disease. *JAMA*. 2016; 316(1):40-50
8. Hauser RA, Factor SA, Marder SR, et al. Kinect 3: A phase 3 randomized, double-blind, placebo-controlled trial of valbenazine for tardive dyskinesia. *American Journal of Psychiatry*. May 2017. 174:5.
9. Waln O, Jankovic J: An update on tardive dyskinesia: from phenomenology treatment. *Tremor Other Hyperkinet Mov (N Y)* 2013; 3: tre-03-161-4138-1.
10. Tarsy, D. Tardive dyskinesia: Prevention and treatment. In: UpToDate, Hurtig, H (Ed). UpToDate, Waltham, MA, 2017.
11. Bhidayasiri, R, Fahn, S, Weiner, WJ, Gronseth, GS, Sullivan, KL, Zesiewicz, TA. Evidence-based guideline: treatment of tardive syndromes: report of the Guideline Development Subcommittee of the American Academy of Neurology. *Neurology*. 2013 Jul 30;81(5):463-9. PMID: 23897874
12. Huntington Study Group, (Marshall F, primary author). Tetrabenazine as antichorea therapy in Huntington disease: A randomized controlled trial. *Neurology* 2006;66:366-372.
13. Micromedex (electronic version, updated periodically). Greenwood Village, CO, USA: Truven Health Analytics.
14. Fasano, A, Bove, F, Lang, AE. The treatment of dystonic tremor: a systematic review. *J Neurol Neurosurg Psychiatry*. 2014;85:759-69. PMID: 24167042