



Dupilumab (Dupixent)

Please fax this completed form to (833) 645-2734 OR mail to: Pharmacy Services | 5 River Park Place East, Suite 210 | Fresno, CA 93720. You can also complete online at [CoverMyMeds.com](https://covermymeds.com).

Coordinated Care of Washington, Inc. Preferred Drug list: https://pharmacy.envolvehealth.com/content/dam/centene/envolve-pharmacy-solutions/pdfs/PDL/FORMULARY-CoordinatedCare_Washington.pdf

Date of request:	Reference #:	MAS:	
Patient	Date of birth	ProviderOne ID or Coordinated Care ID	
Pharmacy name	Pharmacy NPI	Telephone number	Fax number
Prescriber	Prescriber NPI	Telephone number	Fax number
Medication and strength		Directions for use	Qty/Days supply

1. Indicate patient diagnosis:
- Moderate to Severe chronic atopic dermatitis
 - Oral corticosteroid dependent asthma
 - Other. Specify:
 - Asthma with an eosinophilic phenotype
 - Chronic rhinosinusitis with bilateral nasal polyposis
2. Is this prescribed by or in consultation with any of the following (check all that apply):
- Allergy/ Immunology
 - Pulmonology
 - Dermatology
 - Other. Specify:
 - Ear, nose, or throat specialist

For diagnosis of Atopic Dermatitis, complete the following:

3. What is patient's current weight? _____ kg Date taken:

Continuation of therapy for atopic dermatitis:

4. Is there clinical documentation of disease stability or improvement defined by any of the following (check all that apply)?
- At least 20% reduction in body surface area (BSA) involvement
 - Achieved or maintained clear or minimal disease from baseline (equivalent to Investigator's Global Assessment (IGA) score of 0 or 1)
 - Experienced or maintained a decrease in Eczema Area and Severity Index (EASI) score of at least 50%
5. Does patient have documentation of improvement in functional impairment for any of the following? (Check all that apply).
- Improvement in of limitation of activities of daily living (ADLs)
 - Sleep disturbances
 - Skin infections
 - Other. Specify:

New start for atopic dermatitis:

6. Does patient have any of the following (check all that apply)?
- At least 10% body surface area (BSA) involvement
 - A disease severity scale scoring demonstrating severe chronic atopic dermatitis (e.g., Investigator's Global Assessment (IGA) score of 3 or greater; Eczema Area and Severity Index (EASI), Patient Oriented Eczema Measure (POEM); etc.)
 - None of the above

7. Does patient have documentation of functional impairment for any of the following? (Check all that apply)
- Limitation of activities of daily living (ADLs) Skin infections
 Sleep disturbances Other. Specify:
8. Indicate if the patient has a history of failure, intolerance, or contraindication to any of the following (check all that apply):
- For children and adolescents:** Two preferred medium potency topical corticosteroids in the previous 6 months with at least a 28-day trial for each
- For adults:** Two preferred high or very high potency topical corticosteroids in the previous 6 months with at least a 28-day trial for each
- Contraindication(s) to all preferred topical corticosteroids.
- Treatment of sensitive areas (face, anogenital, skin folds) not responding to low potency desonide or hydrocortisone
- History of steroid induced atrophy
- Long-term uninterrupted use
- One preferred topical calcineurin inhibitors (i.e., pimecrolimus, tacrolimus) for daily treatment for at least 28 days
- Contraindication(s) to topical calcineurin inhibitors (i.e., pimecrolimus, tacrolimus)
- Patient age less than 2 years of age
- Phototherapy
- Systemic Immunosuppressants: (i.e., methotrexate, cyclosporine, azathioprine, or mycophenolate)
- Systemic corticosteroids
- Crisaborole (Eucrisa) daily treatment for at least 28 days

For diagnosis of Asthma, complete the following:

Continuation of therapy for asthma with an eosinophilic phenotype or asthma with oral corticosteroid dependent asthma:

9. Is there documentation of disease improvement compared to baseline measures (e.g., reduced missed days from work or school, improved FEV₁, ACQ or ACT scores, decrease in burst of systemic corticosteroids, etc.)?
- Yes No
10. **For asthma with oral corticosteroid dependent asthma:** Has the patient had a reduction in daily oral corticosteroid dosage or usage? Yes No

New start for asthma with an eosinophilic phenotype or asthma with oral corticosteroid dependent asthma:

11. Has patient had any of following (check all that apply):
- FEV₁ less than (<) 80% predicted
- Two or more bursts of systemic corticosteroids in last 12 months
- Poor symptom control (ACQ score consistently greater than 1.5 or ACT score consistently less than 20)
- For diagnosis of asthma with an eosinophilic phenotype:** Frequent (at least twice per year) additional medical treatment such as: emergency department (ED) visits, hospitalizations, treatment with mechanical ventilation, or unplanned (sick) office visits
- For diagnosis of asthma with an eosinophilic phenotype:** Limitation of activities of daily living, nighttime awakening, or dyspnea
12. Will this be used in combination with other monoclonal antibodies (benralizumab, omalizumab, mepolizumab, reslizumab)? Yes No
13. Will patient be using in combination with additional asthma controller medications?
- Yes, please indicate the medication and duration of use. _____

No, please explain. _____

14. **For diagnosis of asthma with oral corticosteroid dependent asthma:** Does the patient have a history of failure (remains symptomatic after 6 weeks) with daily oral corticosteroids, in addition to high-dose inhaled corticosteroids in combination with additional controller(s)? Yes No
15. **For diagnosis of asthma with an eosinophilic phenotype:** Does the patient have a history of failure (remains symptomatic after 6 weeks), contraindication or intolerance to high-dose inhaled corticosteroids, in combination with additional controller(s)? Yes No
16. **For diagnosis of asthma with an eosinophilic phenotype:**
What is patient's blood eosinophil count? _____ cells/ μ L Date taken: _____
17. **For diagnosis of asthma with an eosinophilic phenotype:** Does the patient have a history of failure, contraindication or intolerance to the preferred asthma monoclonal antibodies listed on the AHPDL? Yes No

For diagnosis of chronic rhinosinusitis with nasal polyposis, complete the following:

Continuation of therapy for chronic rhinosinusitis with nasal polyposis:

18. Does patient have clinical documentation of disease improvement compared to baseline defined as a reduction in sinusitis-related symptoms, (such as nasal obstruction, nasal discharge, nasal polyp size, facial pain, and pressure, etc.)? Yes No
19. Will the patient continue to use intranasal corticosteroids with dupilumab? Yes No

New start chronic rhinosinusitis with nasal polyposis:

20. Is there clinical documentation in the patient's file confirming the diagnosis of chronic rhinosinusitis with nasal polyposis? Yes No
21. Does patient have a history of persistent symptoms of rhinosinusitis after completion of 2 months of intranasal corticosteroid use? Yes No
22. Does patient have a history of failure, intolerance, or contraindication to short courses of systemic oral corticosteroids? Yes No
23. Will the patient continue to use intranasal corticosteroids with dupilumab? Yes No

CHART NOTES ARE REQUIRED WITH THIS REQUEST

Prescriber signature	Prescriber specialty	Date
----------------------	----------------------	------

Pharmacy Services will respond via fax or phone within 24 hours of receipt of the request. Requests for prior authorization must include member name, ID#, and drug name. Please include lab reports with requests when appropriate (e.g., Culture and Sensitivity; Hemoglobin A1C; Serum Creatinine; CD4; Hematocrit; WBC, etc.)